

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:
Robert De Leys
Bart Vanderborght
Eric Saman
Hugo Van Heuverswyn

Group Art Unit:

Examiner:

Atty. Dkt. No.: 11362.0025.DVUS03
INNS025--3

Serial No.:


Filed: May 9, 2001

For: PROCESS FOR DETECTING HIV-3
RETROVIRUS (Amended)

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, DC 20231

Sir:

<p style="text-align: center;">CERTIFICATE OF EXPRESS MAIL</p> <p>NUMBER EL521279522US</p> <p>DATE OF DEPOSIT May 9, 2001</p> <p>I hereby certify that this paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to: Commissioner for Patents, Washington, D.C. 20231.</p> <p style="text-align: center;"> Signature</p>

Please amend this application as follows:

IN THE ABSTRACT

Please add the following abstract on a separate page:

--Described is a new variety of retrovirus designated HIV-3, also known as HIV-1 subtype O, samples of which are deposited in the European Collection of Animal Cell Cultures (ECACC) under V88060301. Further described is a process to detect the HIV-3 retrovirus in biological liquids or tissue. One such process involves contacting a biological sample suspected of containing HIV-3 nucleic acids with a DNA probe corresponding to a segment of genomic HIV-3 retrovirus RNA, such as the LTR region, and detecting the hybridization products thereof.--

IN THE SPECIFICATION

On the title page, please delete the original title and insert a new title:

--PROCESS FOR DETECTING HIV-3 RETROVIRUS--

At page 1, line 1, insert a new first paragraph:

--This is a continuation of co-pending application Serial No. 09/379,270, filed August 23, 1999, which is a continuation of 08/900,902, filed July 25, 1997, now issued as United States Patent No. 6,013,484, which is a divisional of Serial No. 08/486,836, filed June 7, 1995, now issued as United States Patent No. 5,795,743, which is a divisional of Serial No. 08/228,519, filed April 15, 1994, now issued as United States Patent No. 5,567,603, which is a divisional of Serial No. 07/460,913, filed March 23, 1990, now issued as United States Patent No. 5,304,466, which claims benefit under 35 U.S.C. §120 of PCT/EP89/00643, filed June 8, 1989 and claims priority under 35 U.S.C. §119 of EP 88 109 200.1, filed June 9, 1988.--

At page 3, line 9, please insert a new paragraph:

--Subsequent to the filing of prior application Serial No. 08/228,519, the medical industry and scientific community has recognized the change in classification of HIV-3 to HIV-1 subtype 0. *See, e.g.,* Rayfield et al., *Emerging Infectious Diseases* 2:209-212 (1996); Janssens et al., *AIDS* 8:1012-1013 (1994); Simon et al., *AIDS* 8:1628-1629 (1994); Gürtler et al., *Journal of Virology* 68:1581-1585 (1994); and Vanden Haesevelde et al., *Journal of Virology* 68:1586-1596 (1994).--

On page 5, lines 24 – 33, please delete the two paragraphs and insert:

--Differential antigen capturing is performed as described hereinafter. The solid line represents the results obtained using a broad-spectrum anti-HIV-1 IgG while the broken line depicts the results obtained using an IgG which was rather specific for HIV-1. Figure 2A, Figure 2B, Figure 2C, Figure 2D, and Figure 2E each shows a typical titration obtained with HIV-1. Figure 2F shows the result obtained with HIV-3 (ANT 70) containing supernatant.

Figure 3A shows differential antigen capturing on HIV-1 and Figure 3B shows differential antigen capturing on HIV03 (ANT 70 NA) supernatants.--

On page 6, lines 7-8, please delete the paragraph and insert:

--Figure 4A shows the reactivity of anti-HIV sera on HIV-1 and Figure 4B shows the reactivity of anti-HIV sera on HIV-2 Western Blot strips.--

On page 7, lines 17-18, please delete the paragraph and insert:

--Figure 8A, Figure 8B, Figure 8C and Figure 8D show the effect of coating IgG dilution on the binding of HIV isolates.--.

On page 8, lines 6-7, please delete the paragraph and insert:

--Figures 10A-E show comparisons of reactivity of human anti-HIV antisera to different HIV types.--

On page 8, lines 19-30, please delete the three paragraphs and insert:

--Figures 11A-C show the titrations of anti-HIV sera by enzyme immunoassay.

Microwell plates were coated with lysates of HIV-1 (SF4), HIV-3 (ANT 70) and HIV-2 (isolate 53). Serum from an HIV-1-infected European (Fig. 11A), antiserum to HIV-3 (ANT 70 NA) (Fig. 11B) and antiserum to HIV-2 (isolate 53) (Fig. 11C) were titrated in 2-fold dilutions beginning at a dilution of 1:100 on all three coated plants.

Figures 12A and B show the positions of methionine and tryptophan residues in viral p17 and p24 gag gene products and Figure 12C shows the positions of methionine and tryptophan residues in viral pol gene products.--

On page 9, lines 11-25, please delete the three paragraphs and insert:

--Figures 13A-D show comparisons of partial cleavage products of gag and pol gene products of HIV-1 (SF4) [HIV-1 in the figure], HIV-3 (ANT 70) [isolate 70 in the figure], HIV-2rod [HIV-2 (LAV-2) in the figure] and HIV-2 (isolate 53) [isolate 53 in the figure]. The terms p24 and p17 are used in the genetic sense to indicate the largest and second largest viral core proteins, respectively.

Figures 14A-1, Figure 14A-II, Figure 14A-III, Figure 14B-1, Figure 14B-II, and Figure 14B-III show hybridization of cDNA clones to viral RNA.

Viral RNA from HIV-1 (SF4), HIV-2rod, and HIV-3 (ANT 70) were spotted onto a membrane filter as described in Materials and Methods. The filters were hybridized under either nonstringent (A) or stringent conditions (B) and autoradiographed.--

At page 55, lines 33- 37 please delete the paragraph and insert:

--The hybridization data also support the notion that ANT 70 is fundamentally different from either HIV-1 and HIV-2. As long as the conditions under which the hybridization is performed are stringent, a distinction can easily be made among the three virus types. RNA of the HIV-3 retrovirus virtually hybridizes neither with the Env gene or the LTR close to it, in particular not with the nucleotide sequence 8352-9538 of HIV-1, nor with the sequences of the Pol region of the HIV-1 genome under stringent conditions.--

IN THE CLAIMS

Cancel claims 1-30 and 32-36, without prejudice.

Please amend claim 31 to read:

31. (Amended) A process for the detection of HIV-3 retrovirus or of its RNA in a biological liquid or tissue, characterized by contacting nucleic acids contained in said biological liquid or tissue with a DNA probe containing at least 360 contiguous sequences corresponding to the genomic RNA of HIV-3 retrovirus under stringent hybridization conditions, washing the hybrid formed with a solution preserving said stringent conditions, and detecting the hybrid formed.

Please add claims 37-40 as follows:

--37. The process of claim 31 wherein the DNA probe is:

10 CCCATGGATT	20 TGAAGATACA	30 CATAAAGAAA	40 TACTGATGTG	50 GAAGTTTGAT	60 AGATCTCTAG
70 GCAACACCCA	80 TGTTGCTATG	90 ATAACTCACC	100 CAGAGCTCTT	110 CCAGAAGGAC	120 TAAAAACTGC
130 TGACCTGAAG	140 ATTGCTGACA	150 CTGTGGAACT	160 TTCCAGCAAA	170 GACTGCTGAC	180 ACTGCGGGGA
190 CTTCCAGTG	200 GGAGGGACAG	210 GGGGCGGTTC	220 GGGGAGTGGC	230 TAACCCTCAG	240 AAGCTGCATA
250 TAAGCAGCCG	260 CTTCTGCTT	270 GTACCGGGTC	280 TCGGTTAGAG	290 GACCAGGTCT	300 GAGCCCGGGA
310 GCTCCCTGGC	320 CTCTAGCTGA	330 ACCCGCTCGT	340 TAACGCTCAA	350 TAAAGCTTGC	360 CTTGAGTGAG

A

or the complement thereof.

38. The process of claim 31 wherein the DNA probe is:

10 AACATGGGAA	20 ACGCATTGAG	30 AAAAGGTAAA	40 TTTGAGGGAT	50 GGGCAGCAGT	60 AAGAGAAAGA
70 ATGAGAAGAA	80 CTAGAACTTT	90 CCCTGAGTCT	100 GAACCATGCG	110 CACCTGGAGT	120 AGGACAGATC
130 TCCAGGGAAT	140 TAGCAGCTAG	150 AGGAGGGATA	160 CCAAGTTCCC	170 ATACTCCTCA	180 AAACAATGCA
190 GCCCTTG CAT	200 TCCTAGAAAG	210 TCACCAAGAG	220 GAAGAAGTAG	230 GTTTTCCAGT	240 AGCACCTCAA
250 GTGCCTCTAA	260 GGCCAATGAC	270 CTATAAAGGA	280 GCATTTGACC	290 TCAGCTTCTT	300 TTTAAAAGAA
310 AAGGGAGGAC	320 TGGAAGGGTT	330 AATTTACTCC	340 CATAAAAGAG	350 CAGAAATCCT	360 GGATCTTTGG

GTGTATAA

or the complement thereof.

39. The process of claim 31 wherein the DNA probe corresponds to the nucleotide sequence coding for proteins p12, p16 or p25 of the HIV-3 retrovirus or the complement thereof.

40. The process of claim 31 wherein the DNA probe corresponds to the nucleotide sequence coding for glycoproteins gp41 or gp120 of the HIV-3 retrovirus or the complement thereof.--

REMARKS

The active claims in this case are claims 31 and 37-40.

The specification has been amended to recite the relationship with the parent cases, and to incorporate portions of the preliminary amendment filed August 23, 1999 in parent case 09/379,270. A marked up version of the amendments to the Specification is attached hereto. The original application was filed without an abstract of the disclosure. This Preliminary Amendment is being filed to provide such an abstract. The title has been amended to reflect the subject of the present claims.

The specification at page 3 has been amended to introduce the current classification, HIV-1 subtype O virus, for what had been termed "HIV-3" in the original application. After the inventors first reported on their discovery of HIV-3, specifically variant ANT₇₀, the medical and scientific community recognized that HIV-3 should more appropriately be classified as a subtype of HIV-1. This subtype was designated "O" where O stands for "outliers". Several journal articles have been provided in the preceding case Serial No. 08/486,836 to substantiate the scientific recognition that HIV-3, e.g., ANT₇₀, is now classified as HIV-1 subtype O.

The new claims find support at page 30, line 25 through page 32, line 16; pages 37-40; pages 50-53; and original claims 25-31. A marked up version of the claim amendments is attached.

It is believed that no fee is due; however, should any fees under 37 C.F.R. §§ 1.16 to 1.21 be required for any reason, the Commissioner is authorized to deduct said fees from Deposit Account No. 01-2508/11362.0025.DVUS03.

Respectfully submitted,



Patricia A. Kammerer

Reg. No. 29,775

Attorney for Assignee
INNOGENETICS N.V.

HOWREY SIMON ARNOLD & WHITE LLP
750 Bering Drive
Houston, TX 77057
(713) 787-1400

Date: May 9, 2001

MARKED-UP VERSION OF CLAIMS

31. (Amended) A process for the detection of HIV-3 retrovirus or of its RNA in a biological liquid or tissue, characterized by contacting nucleic acids contained in said biological liquid or tissue with a DNA probe containing [a nucleic acid according to any of claims 25 to 30] at least 360 contiguous sequences corresponding to the genomic RNA of HIV-3 retrovirus under stringent hybridization conditions, washing the hybrid formed with a solution preserving said stringent conditions, and detecting the hybrid formed.

--37. The process of claim 31 wherein the DNA probe is:

10 CCCATGGATT	20 TGAAGATACA	30 CATAAAGAAA	40 TACTGATGTG	50 GAAGTTTGAT	60 AGATCTCTAG
70 GCAACACCCA	80 TGTTGCTATG	90 ATAACTCACC	100 CAGAGCTCTT	110 CCAGAAGGAC	120 TAAAAACTGC
130 TGACCTGAAG	140 ATTGCTGACA	150 CTGTGGAAC	160 TTCCAGCAAA	170 GACTGCTGAC	180 ACTGCGGGGA
190 CTTTCCAGTG	200 GGAGGGACAG	210 GGGGCGGTTC	220 GGGGAGTGGC	230 TAACCCTCAG	240 AAGCTGCATA
250 TAAGCAGCCG	260 CTTCTGCTT	270 GTACCGGGTC	280 TCGGTTAGAG	290 GACCAGGTCT	300 GAGCCCGGGA
310 GCTCCCTGGC	320 CTCTAGCTGA	330 ACCCGCTCGT	340 TAACGCTCAA	350 TAAAGCTTGC	360 CTTGAGTGAG

A

or the complement thereof.

38. The process of claim 31 wherein the DNA probe is:

10 AACATGGGAA	20 ACGCATTGAG	30 AAAAGGTAA	40 TTTGAGGGAT	50 GGGCAGCAGT	60 AAGAGAAAGA
70 ATGAGAAGAA	80 CTAGAACTTT	90 CCCTGAGTCT	100 GAACCATGCG	110 CACCTGGAGT	120 AGGACAGATC
130 TCCAGGGAAT	140 TAGCAGCTAG	150 AGGAGGGATA	160 CCAAGTTCCC	170 ATACTCCTCA	180 AAACAATGCA
190 GCCCTTGCA	200 TCCTAGAAAG	210 TCACCAAGAG	220 GAAGAAGTAG	230 GTTTTCCAGT	240 AGCACCTCAA
250 GTGCCTCTAA	260 GGCCAATGAC	270 CTATAAAGGA	280 GCATTTGACC	290 TCAGCTTCTT	300 TTTAAAAGAA

310 320 330 340 350 360
AAGGGAGGAC TGGAAGGGTT AATTTACTCC CATAAAAGAG CAGAAATCCT GGATCTTTGG
GTGTATAA
or the complement thereof.

39. The process of claim 31 wherein the DNA probe corresponds to the nucleotide sequence coding for proteins p12, p16 or p25 of the HIV-3 retrovirus or the complement thereof.

40. The process of claim 31 wherein the DNA probe corresponds to the nucleotide sequence coding for glycoproteins gp41 or gp120 of the HIV-3 retrovirus or the complement thereof.--

"EXPRESS MAIL" MAILING LABEL

NUMBER EM 219636253 US

DATE OF DEPOSIT July 25, 1997

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS
BEING DEPOSITED WITH THE UNITED STATES POSTAL
SERVICE, "EXPRESS MAIL POST OFFICE TO ADDRESSEE"
SERVICE UNDER 37 C.F.R. 1.10 ON THE DATE INDICATED
ABOVE IS ADDRESSED TO THE ASSISTANT COMMISSIONER
FOR PATENTS, WASHINGTON, D.C. 20231.

**HIV-3 RETROVIRUS AND ITS USE
PROCESS FOR DETECTING**

Inventors: Robert De Leys

Bart Vanderborcht

Eric Saman

Hugo Van Heuverswyn

"EXPRESS MAIL" MAILING LABEL

NUMBER IB 853220214

DATE OF DEPOSIT 6-7-95

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS
BEING DEPOSITED WITH THE UNITED STATES POSTAL
SERVICE, "EXPRESS MAIL POST OFFICE TO ADDRESSEE"
SERVICE UNDER 37 C.F.R. 1.10 ON THE DATE INDICATED
ABOVE IS ADDRESSED TO THE COMMISSIONER OF
PATENTS AND TRADEMARKS, WASHINGTON, D.C. 20231.

Maria L. Alvarez

"EXPRESS MAIL" MAILING LABEL

NUMBER NB 162 253 168

DATE OF DEPOSIT February 9, 1990

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS
BEING DEPOSITED WITH THE UNITED STATES POSTAL
SERVICE, "EXPRESS MAIL POST OFFICE TO ADDRESSEE"
SERVICE UNDER 37 C.F.R. 1.10 ON THE DATE INDICATED
ABOVE AND IS ADDRESSED TO THE COMMISSIONER OF
PATENTS AND TRADEMARKS, WASHINGTON, D.C. 20231.

Mary A. Paul
SIG

<p>"EXPRESS MAIL"</p> <p>Number <u>RB 700732037</u></p> <p>Date of Deposit <u>4-15-94</u></p> <p><small>Pursuant to 37 C.F.R. § 1.10, I hereby certify that I am personally depositing this paper or fee with the U.S. Postal Service, "Express Mail Post Office to Addressee" service on the date indicated above in a sealed envelope (a) having the above-numbered Express Mail label and sufficient postage affixed, and (b) addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.</small></p> <p><u>John Dargatzis</u> Signature</p>

1

source of virus.

5

Furthermore, the invention relates to a purified retrovirus having the essential morphological and immunological properties described below. In many cases, the unique characteristics of HIV-3 can best be appreciated by comparison with the same type of characteristics relating to the other human immunodeficiency viruses, HIV-1 and HIV-2.

10

Brief description of the drawings

15

In Figures 1 to 16 the designations HIV-3 (ANT 70) and HIV-3 (ANT 70 NA) refer to two strains of a new HIV-3 virus isolated from a Camerounian woman and her partner from which HIV-3 (ANT 70) has been deposited under ECACC V88060301.

20

Figure 1 shows a procedure for preparing cleavage maps of viral proteins.

25

Figure 2 shows differential antigen capturing on virus-containing culture supernatants.

Differential antigen capturing is performed as described hereinafter. The solid line represents the results obtained using a broad-spectrum anti-HIV-1 IgG while the broken line depicts the results obtained using an IgG which was rather specific for HIV-1. ~~The titrations shown in panels A-E are typical for HIV-1. Panel F shows the result obtained with HIV-3 (ANT 70) containing supernatant.~~

30

Figure 3A shows differential antigen capturing on HIV-1 and HIV-3 (ANT 70 NA) supernatants.

35

Figure 3B shows differential antigen capturing on Differential antigen capturing was performed as described hereinafter. The solid line depicts the results obtained on

Figure 2A, Figure 2B, Figure 2C, Figure 2D, and Figure 2E each show a typical titration obtained with HIV-1. Figure 2F

1

plates coated with the broad spectrum anti-HIV IgG while the broken line represents the results obtained on plates coated with IgG which shows less crossreactivity with HIV types other than HIV-1.

5

Figure 4 shows the reactivity of anti-HIV sera on HIV-1 and HIV-2 Western Blot strips.

↑ *Figure 4B shows the reactivity of anti-HIV sera on*

10 The reactivities of 3 different sera on HIV-1 and HIV-2 Western blot strips are shown. Sera: 1. anti-HIV-1, 2. anti-HIV-3 (ANT 70), 3. anti-HIV-2 (isolate 53). The molecular weights indicated are those given by the manufacturer (Dupont Biotech).

15

Figure 5 relates to the comparison of gag and pol proteins of several HIV-1 isolates, HIV-2rod and HIV-3 (ANT 70).

20

Proteins were separated electrophoretically and blotted as described later. The blot was incubated with a broad-spectrum anti-HIV antiserum followed by (anti-human IgG)/alkaline phosphatase- labeled conjugate to visualize the proteins.

25

A. HIV-2rod, B. an HIV-1 laboratory isolate, C. HIV-3 (ANT 70), D. an HIV-1 laboratory isolate, E. HIV-1 (SF4).

Figure 6 shows a comparison of HIV-3 (ANT 70) and HIV-3 (ANT 70 NA) proteins.

30

Proteins were separated electrophoretically and blotted as described later. The blot was incubated with the BSR antiserum followed by (alkaline phosphatase)/anti-human IgG conjugate to visualize the proteins. Lane 1: HIV-3 (ANT 70 NA), lane 2: HIV-3 (ANT 70), lane 3: HIV-1 (SF4). The apparent intensity difference between lanes 1 and 2 is caused by the difference in the amount of material loaded.

35

1

Figure 7 relates to the ability of various human anti-HIV-1 sera to capture viral antigens.

5

A number of human sera were diluted 1:1000 and coated directly on microwell plates. Detergent-treated culture supernatants containing HIV-1 (SF4), HIV-3 (ANT 70), HIV-2rod or HIV-2 (isolate 53) were incubated and the bound antigen was detected using a broadspectrum (anti-HIV)/horseradish peroxidase conjugate. Sera 1-7 were of African origin while sera 8-11 were from Europeans. The greater ability of African sera to capture non-HIV-1 antigen can, in part, be explained by their higher anti-p24 titers (data not shown).

15

A, Figure 8B, Figure 8C and Figure 8D show

Figure 8 shows the effect of coating IgG dilution on the binding of HIV isolates.

20

Successive 2-fold dilutions were made of four different sera, beginning at a dilution of 1:1000 and were used to coat microwell plates. Detergent-treated supernatants of HIV-1 (SF4), HIV-3 (ANT 70), HIV-2rod and HIV-2 (isolate 53) were diluted to give approximately the same optical density on plates coated with the antiserum shown in panel B at a dilution of 1:1000. Bound antigen was detected using the broad-spectrum (anti-HIV IgG)/horseradish peroxidase conjugate.

25

30

Figure 9 shows antigen capturing of virus isolates using human polyclonal and mouse anti-HIV-1 monoclonal antibodies.

Wells were coated and incubated as described in the text. The IgGs used are as follows:

35

1. human polyclonal anti-HIV IgG , 2. MAb CLB 59, 3. MAb CLB

21, 4. MAb CLB 64, 5. MAb CLB 14, 6. MAb CLB 16, 7. MAb CLB 47, 8. MAb CLB 13.6 (anti-p18), 9. MAb CLB 19.7, 10. MAb CLB 13.4 (anti-p18).

5 *A-E show comparisons*
~~Figure 10 is a comparison~~ of the reactivity of human anti-HIV antisera to different HIV types.

10 Lysates of HIV-1 (SF4), HIV-3 (ANT 70), HIV-2rod and HIV-2 (isolate 53) were separated electrophoretically on SDS-polyacrylamide gels, blotted onto nitrocellulose, and incubated with a high titer anti-HIV-1 antiserum (panel A), a lower titer anti-HIV-1 antiserum (panel B), serum from the woman from whom HIV-3 (ANT 70) was isolated (panel C), her partner from which HIV-3 (ANT 70 NA) was isolated (panel D) and anti-HIV-2 antiserum from the person from whom HIV-2 (isolate 53) was isolated (panel E).

15 *A-C show*
~~Figure 11 shows~~ titrations of anti-HIV sera by enzyme immunoassay.

20 Microwell plates were coated with lysates of HIV-1 (SF4), HIV-3 (ANT 70) and HIV-2 (isolate 53). Serum from an HIV-1-infected European (*Fig. 11A* left panel), antiserum to HIV-3 (*Fig. 11B* center panel) and antiserum to HIV-2 (isolate 53) (*Fig. 11C* right panel) were titrated in 2-fold dilutions beginning at a dilution of 1:100 on all three coated plates.

25 *A and B show*
~~Figure 12 shows~~ the positions of methionine and tryptophan residues in viral gag and pol gene products.

p17 and p24 gene products
Amino acid positions for the p17 gag proteins are given starting from the first methionine in the coding sequence. *And Figure 12C shows the positions of methionine and tryptophan residues in viral*
Positions for the p24 gag protein are given starting at the p17/p24 proteolytic cleavage site. Positions for the pol gene are shown after alignment with the highly conserved

1

tryptophan doublet in the HIV-1 sequence at positions 556 and 557. The positions of a conserved protease sequence, the protease/reverse transcriptase cleavage site and the reverse transcriptase/endonuclease cleavage site are indicated. In this case, the terms p24 and p17 are used in the genetic sense to refer to the largest and second largest viral core proteins respectively. The term "HIV-2 (LAV-2)" is a synonymum for HIV-2 rod.

5

10

~~Figures 13A-D show comparisons.~~

~~Figure 13 is a comparison~~ of partial cleavage products of gag and pol gene products of HIV-1 (SF4) [HIV-1 in the figure], HIV-3 (ANT 70) [isolate 70 in the figure], HIV-2rod [HIV-2 (LAV-2) in the figure] and HIV-2 (isolate 53) [isolate 53 in the figure]. The terms p24 and p17 are used in the genetic sense to indicate the largest and second largest viral core proteins, respectively.

15

~~Figures 14A-I, Figure 14A-II, Figure 14A-III, Figure 14B-I, Figure 14B-II and Figure 14B-III show~~
~~Figure 14 shows~~ hybridization of cDNA probes to viral RNA.

20

Viral RNA from HIV-1 (SF4), HIV-2rod, and HIV-3 (ANT 70) were spotted onto a membrane filter as described in Materials and Methods. The filters were hybridized under either nonstringent (A) or stringent conditions and autoradiographed.

25

(B)

1. Morphology

30

Electron microscopy of HIV-3-infected MT4 cells revealed the presence of extracellular virus particles having a diameter

35

1 antigenic differences between ANT 70 and HIV-1 are
smaller than those between HIV-2 and HIV-1. This is
particularly evident from the results presented in Figures 8
and 10.

5

Additional compelling evidence that ANT 70 is a unique
virus different from HIV-1 and HIV-2 comes from the partial
peptide maps. We have shown that there are significant
differences in the most highly conserved viral proteins.

10 The two HIV-2 isolates which were used for comparison gave
essentially identical cleavage patterns except in the case
of CNBr cleavage of the p17 core protein. It should be
noted, however, that the p17 core protein exhibits more
variability than the p24 protein, at least in HIV-1 strains
15 (34). Whether or not this also holds true for HIV-2 awaits
sequence determination on more strains than have been
analyzed to date.

In light of the fact that ANT 70 is antigenically more
closely related to HIV-1 than is HIV-2, as evidenced by a
20 higher degree of crossreactivity which extends even to the
gp41 envelope protein, was essential to establish that
ANT 70 was more than simply a genetic variant of HIV-1.
This was possible by investigating the locations of some of
the most highly conserved amino acids in a number of viral
25 proteins which are least subject to genetic variation. That
major differences were noted in the cleavage patterns
indicates that HIV-1, HIV-2 and ANT 70 are three genetically
distinct viruses. On the other hand, the same series of
experiments also revealed similarities between the viruses
30 which may indicate that all three arose from a common
progenitor.

The hybridization data also support the notion that ANT 70
is fundamentally different from either HIV-1 and HIV-2. As
35 long as the conditions under which the hybridization is
performed are stringent, a distinction can easily be made
among between the three virus types. RNA of the HIV-3
retrovirus virtually hybridizes neither with the Env
gene or the LTR close to it, in particular not
with the nucleotide sequence 8352-9538 of HIV-1,
nor with the sequences of the Pol region of the HIV-1
genome under stringent conditions.